Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 1 January 2013 (MN LaTeX style file v2.2)

arXiv:1212.6630v1 [astro-ph.GA] 29 Dec 2012

Detailed studies of the PSR B1259−63 spectrum evolution and classification of radio pulsar spectra

M. Dembska⋆, J. Kijak and W. Lewandowski
Kepler Institute of Astronomy, University of Zielona Góra, Lubuska 2, 65-265 Zielona Góra, Poland

Accepted... Received...; in original form...

ABSTRACT

Kijak et al. (2011a) studied the radio spectrum of PSR B1259−63 in an unique binary with Be star LS 2883 and showed that this pulsar undergoes a spectrum evolution due to orbital motion. They proposed a qualitative model which explains this evolution. They considered two mechanisms that might influence the observed radio emission: free-free absorption and cyclotron resonance. Using the same database we constructed spectra for chosen observing days and obtained different types of spectra. Comparing to current classification of pulsar spectra, there occurs a suggestion that the appearance of various spectra shapes, different from a simple power law which is typical for radio pulsars, is possibly caused by environmental conditions around neutron stars. Therefore, the case of B1259−63 can be treated as a key factor to explain not only the GPS phenomenon observed for the solitary pulsars with interesting environments and also another types of spectra (e.g. with break).

Key words: pulsars: general, individual (B1259−63) - stars: winds, outflows - ISM: general

1 INTRODUCTION

The shapes of pulsar spectra are very important in understanding of the pulsar radio emission. Typical radio spectrum of a pulsar can be described using either a simple power law with a negative spectral index of -1.8 or two power laws with spectral indices of -0.9 and -2.2 and a break frequency on average of 1.5 GHz (Maron et al. 2000). For several pulsars there were observed a low frequency turnover (Sieber 1973; Malofeev et al. 1994). Lorimer et al. (1995) also mentioned about some pulsars having positive spectral index or flat spectrum in the frequency range between 400 MHz and 1600 MHz.

Kijak et al. (2011b) provided a definite evidence for a new type of pulsar radio spectra. These spectra show the maximum flux above 1 GHz, while at higher frequencies the spectra look like a typical pulsar spectrum. At lower frequencies (below 1 GHz), the observed flux decreases, showing a positive spectral index (Kijak et al. 2011b). They called these objects the gigahertz-peaked spectra (GPS) pulsars. A frequency at which such a spectrum shows the maximum flux was called the peak frequency. Kijak et al. (2011b) also indicated that the GPS pulsars are relatively young objects and they usually adjoin such interesting environments as HII regions or compact pulsar wind nebulae (PWN). Additionally some of them seem to be coincident with the known but sometimes unidentified X-ray sources from 3rd EGRET Catalogue or HESS observations.

Candidates for GPS pulsars were selected by Kijak & Maron (2004) amongst those which showed a decreasing flux density at frequencies below 1 GHz. Kijak et al. (2007) used multifrequency flux measurements for these candidate pulsars and presented the first direct evidence of a high frequency turnover. All of them have proved to be relatively young and to have high value of dispersion measure (DM around 150 $\mathrm{pc\,cm^{-3}}$ and above). PSR B1259−63 was also inscribed by Lorimer et al. (1995) in the list of pulsars with positive spectral indices. This pulsar is in an unique binary with a massive main-sequence Be star. B1259−63 has a short period of 48 ms and a characteristic age of 330 kyr which places it amongst middle-aged pulsars. Its average DM is about 147 $\mathrm{pc\,cm^{-3}}$ and the corresponding distance is about 2.75 kpc. Therefore, B1259−63 seems a natural candidate to be classified as the GPS pulsar. The companion star LS 2883 is a 10-mag massive Be star with a mass of about $10\mathrm{M}_{\odot}$ and a radius of $6\mathrm{R}_{\odot}$ (Johnston et al. 1994). Negueruela et al. (2011) estimates the mass of LS 2883 of about $30\mathrm{M}_{\odot}$, however the result is subject to large uncertainties. Be stars are generally believed to have a hot tenuous polar wind and a cooler high-density equatorial disc. The PSR B1259−63/LS 2883 emits unpulsed non-thermal emission over a wide range of frequencies ranging between radio and γ-rays, and its flux varies with orbital phase. Kijak et al. (2011a) studied the radio spectrum of B1259−63 using the available measurements of the pulsed flux obtained during three periastron passages (1997, 2000 and 2004) and showed that this pulsar undergoes a spectrum evolution due to its orbital motion. In addition, demonstrated that the peak frequency also depends on the orbital phase and therefore it varies with the changes of the pulsar environment. They argued that the observed variation of the spectra is caused by a combination of two effects: the free-free absorption in the stellar wind and the cyclotron resonance in the magnetic field. This field is as-

---

⋆ E-mail: marta@astro.ia.uz.zgora.pl

Table 1. The results of our best fit choice to B1259−63 spectra for each observing day, ”-” and ”+” denote days before and after periastron passage respectively, year is the epoch of observation.

Day	Year	Best fit	Comments
-60	1997	turnover	
-48	1997	unclear	no trend
-46	2000	broken	
-40	1997	turnover	
-24	1997	turnover	
-21	2004	turnover	
-18	2000	turnover	
+16	1997	turnover	
+20	1997	turnover	
+21	2004	broken/turnover	small positive spectral index below 2 GHz
+24	2000	turnover	
+27	1997	turnover	
+28	2004	power-law	
+29	2000	power-law	
+30	1997	unclear	no trend
+33	2004	power-law/broken	small break
+34	2000	broken/turnover	small positive spectral index below 4.2 GHz
+36	1997	turnover	
+38	2004	broken/unclear	difficult to classify
+40	2000	unclear	no trend
+42	1997	flat	
+47	1997	broken/turnover	small positive spectral index below 2.5 GHz
+55	2000	unclear	no trend
+63	2000	broken	
+64	2004	turnover	
+70	2000	power-law/broken	small break
+83	2000	broken	
+94	2004	broken	
+113	2000	unclear	significant error bars
+150	2004	broken	
+186	2004	power-law	

sociated with the disc and is infused by the relativistic particles of the pulsar wind. The apparent resemblance between the B1259−63 spectrum and the GPS suggests that the same mechanisms should be responsible for both cases (Kijak et al. 2011a). Thus, there appears a conclusion that the GPS feature should be caused by some external factors rather than by the emission mechanism.

We collected flux density observations of B1259−63 measured during three periastron passages (Johnston et al. 1999; Connors et al. 2002; Johnston et al. 2005) and constructed radio spectra for different observing days. Then we analysed their shapes in details taking into account the influence of interstellar scintillations and expressed implications of B1259−63 case on classification of radio pulsar spectra in general.

2 B1259−63 RADIO SPECTRUM: DETAILED ANALYSIS

As it was mentioned previously, the typical pulsar spectrum is usually described in terms of a simple-power law

$$F(\nu) = c\nu^{\alpha}, \tag{1}$$

except from a frequency range below some critical value where we presume a cut-off which may be caused by a loss of coherence. However, there are also known some more complicated spectra, i.e. turnover or broken-type. In case of spectrum with so-called break the usual next step is fitting two power laws to the inspected data

$$F(\nu) = \begin{cases} c_1\nu^{\alpha_1} : & \nu \leqslant \nu_b \\ c_2\nu^{\alpha_2} : & \nu > \nu_b \end{cases} \tag{2}$$

where ν_b denotes break frequency. Broken-type spectrum is usually defined by having two negative spectral indices and becoming steeper at higher frequencies. Spectrum with turnover, either at low or high frequency, can be modelled in a way proposed by Kuzmin & Losovsky (2001)

$$F(\nu) = 10^{ax^2+bx+c}, \quad x \equiv \log_{10}\nu, \tag{3}$$

and $\nu_p = 10^{\frac{-b}{2a}}$ denotes peak frequency. Kijak et al. (2011a) used the same method to analyse both the shapes of GPS and the spectrum evolution of PSR B1259−63 for chosen intervals of orbital phases and showed that the pulsar undergoes spectrum evolution while it orbits around the companion. It is important to mention that they used data with available four flux measurements for a given observing day except −18d (i.e. 18 days prior to the periastron passage), corresponding to the orbital phase which is just before the eclipse, with only three observing points. We created a database with all available flux density measurements for this pulsar which allowed us to re-examine the spectrum evolution presented by Kijak et al. (2011a). Despite using additional observing points which weren’t been taken into consideration previously, it appears that there wasn’t significant differences in shapes of B1259−63 spectra for chosen orbital phase intervals between our results and those from Kijak et al. (2011a). However, our analysis revealed new implications on classification of radio pulsar spectra.

To clarify the spectrum evolution, let us examine the B1259−63 spectrum observed compiled for each observing day using the same database. We choose methods which is a combination of those used by Maron et al. (2000) for radio spectra of about 280 pulsars and Kijak et al. (2011a) for GPS and B1259−63 spectrum evolution. For reliable results we analyse only spectra with four observing points with additional spectrum for −18d constructed of three flux measurements.

2.1 Data analysis

To analyse the spectra in details, we attempted to fit both one/two power law(s) to the data with errors. Moreover, a simple parabolic function was used instead of this given in equation Eq.(3). We accomplish the fitting using an implementation of the linear least-squares fitting algorithm (available in GNU Scientific Library) to logarithms of observational data. To choose the best fit we had to analyse the results in a complex way. At first, we rejected the evident inaccurate approximations and non-physical results, e.g. break at frequency below the lowest or above the highest one. Taking into account that there are available maximum four data points we could not simply compare the values of χ^2 to denote the best fit. Otherwise, we would classify most of spectra as broken-type as fitting two power law functions to 2-elements data set each is statistically the most accurate one. Let us denote that resting on an existing classification of radio pulsars spectra there are three basic types of spectra.

As it was mentioned, we expect broken-type of spectrum to have two negative spectral indices. This in turn leads us to classify



Figure 1. The fits to the B1259−63 spectra for the each orbital phase interval, prior (left panel) and after (right panel) periastron passage.

Table 2. Best fit and parameters fitted to spectra presented on Fig.2, for more details see Eq. (1), (2) and (3); "-" and "+" denote days before and after periastron passage respectively, year is the epoch of observation.

Day	Year	Best fit	Parameters
+33	2004	power-law	$c = 6.9$, $\alpha = -0.98$
+83	2000	broken	$c_1 = 5.6$, $\alpha_1 = -0.18$, $c_2 = 69.7$, $\alpha_2 = -2.3$, $\nu_b = 3.3$ GHz
+20	1997	turnover	$a = -4.7$, $b = 5.1$, $c = -0.74$, $\nu_p = 3.5$ GHz
+42	1997	flat	$c = 3.8$, $\alpha = -0.15$

spectra showing positive spectral index below some peak frequency as turnover spectra. We also specify power-law spectra with very small value (close to zero) of spectral index as a flat one. Some of spectra showed two different trends (for more details see Tab.1). Any other existing cases would be given recognition as unclear. We present the results of best fit for each spectrum in Table 1.

2.2 Results

Kijak et al. (2011a) presented detailed study of PSR B1259−63 spectra constructed for few orbital epochs. They also analysed published data of the pulsar flux measured during three periastron passages in 1997, 2001 and 2004 but they calculated an average flux density using only those data with observing points at least four given frequencies for a given observing day. We extended the database to include all measurements that have been published so far. Fig.1 shows the spectra for various orbital phase range which can be used as a reference to analysed spectra for given observing days. It seems clear that the shape of the PSR B1259−63 spectrum depend on the orbital phase. Therefore, we can acknowledge that the PSR B1259−63 spectrum undergoes evolution. Kijak et al. (2011a) suggested that free-free absorption due to the stellar wind can be responsible for this effect. Moreover, our analysis confirms that the shape of spectra are not fully symmetric with respect to the periastron point (Kijak et al. 2011a). One can note that flux values on the left panel are generally smaller that those on the right panel of Fig.1. This effect can be explained by the fact that the waves emitted before the periastron passage travel longer distance through the stellar wind (see Fig. 4 in Kijak et al. (2011a)). Thus, they are affected by free-free absorption stronger than the waves emitted after the eclipse. We can also confirm that a peak (or in some cases a break) frequency depend on the orbital phase. This can be cause by the cyclotron resonance in magnetic field of the disc (Kijak et al. 2011a).

Comparing the results of fitting presented in Tab.1 one can note that shapes of spectra for single days are generally consistent with those averaged over orbital phase intervals. There also can be seen some kind of domination of GPS feature for orbital epoch before eclipse. The same time the general scenario leads spectra after eclipse from turnover to other shapes. It is important to mention that first sign of more typical spectrum (i.e. broken or power-law) appears as early as 20 days after periastron passage. The flux at the given frequency apparently changes with orbital phases. When the pulsar is close to periastron, the flux generally decreases at all observed frequencies and the most drastic decrease is observed at the lowest frequency. Moreover, we noticed all types of radio pulsar



Figure 2. The fits to the B1259−63 spectra for chosen days (”-” and ”+” denote days before and after periastron passage respectively). Each panel shows different type of spectra.

spectra, including a flat spectrum (see Fig. 2). Close to the periastron point the spectra of B1259−63 resemble those of the GPS pulsars. The spectrum for the orbital epochs further from the periastron point are more consistent with typical pulsar spectra (i.e. power-law and broken).

3 DISCUSSION

3.1 Interstellar scintillations influence on B1259−63 spectra

Pulsars are generally known to be stable radio sources although the pulsar signal is affected in a number of ways as it propagates through the interstellar medium (Lorimer & Kramer 2005). Thus, the measured flux density varies due to diffractive and refractive scintillation effect. Scintillation studies have been extremely successful in determining some properties of the interstellar medium and the location of the dominant scattering screen. PSR B1259−63 is a object with relatively high dispersion measure (which means that its transition frequency is very high) and at the frequencies at which it was observed it should be in the so called *strong scintillation regime*. This implies that we definitely have to take into consideration both refractive (RISS) and diffractive (DISS) scintillations when analysing spectra for a given day. We used diffractive bandwidth $\Delta f_{\rm DISS}$ and timescale $\Delta t_{\rm DISS}$ from scintillation observations of the pulsar made far from periastron at 4.8 GHz and 8.4 GHz (McClure-Griffiths et al. 1998) to estimate values of these parameters at 1.4 GHz and 2.4 GHz assuming $\Delta t_{\rm DISS} \propto f^{1.2} d^{-0.6}$ (Lorimer & Kramer 2005), where f and d denote frequency and distance respectively. McClure-Griffiths et al. (1998) observed PSR B1259−63 during 46 independent observations between 1993 August and 1997 February, excluding data taken in 1993 December and 1994 January during the periastron passage. The observations were typically between 30 and 100 minutes long. The standard deviation of the diffractive bandwidths and timescales values at 4.8 GHz and 8.4 GHz throughout the orbit is ∼30% (McClure-Griffiths et al. 1998). We estimated the values of $\Delta t_{\rm DISS}$ to be ranging from 40s at 1.4 GHz to 360s at 8.4 GHz which suggests that diffractive scintillations should not affect the average flux measurements (observing sessions was usually 4 hours long). Roughly estimated refractive timescales vary from 12 hours at 8.4 GHz to more than 20 days at 1.4 GHz. However, for lower frequencies the modulation index is relatively small which means lower uncertainty estimates when measuring flux. However, we can definitely point out at least a couple of high frequency observations will be affected by refractive scintillations, e.g. for orbital phase from +113 to +186 days (see Tab. 1) including only three spectra, one of which is unclear. This leads to conclusion that flux values should be averaged over epochs and/or orbital phase intervals to be more reliable. However, it is important to mention that shapes of spectra for single days undergo evolution which is similar in a significant way to this observed for spectra averaged orbital phase intervals.

3.2 Implications of B1259−63 case on classification of radio pulsar spectra

Maron et al. (2000) concluded that the single power law spectrum is the most typical for a sample of about 280 pulsars in a wide frequency range. Their studies showed that only 10% of pulsars requires two power law spectra. Both numbers of low and high frequency turnover spectra are also still relatively small (Sieber 1973; Maron et al. 2000; Kijak et al. 2011a). These results indicate the possible existence of the ”true” shape of spectrum for each pulsar and other types of spectra being a rare exceptions of an unknown origin.

It seems clear that the existing classification of radio pulsar spectra is stricte morphological and any direct connection between external or internal factors and the shape of spectrum cannot be drawn. Kijak et al. (2011a) suggested that the GPS feature should be caused by some external factors rather than by the emission mechanism. One can note that shapes of spectra for single days presented in Tab.1 confirm strong environmental influence on the pulsar radio emission. This leads us to suggestion that occurrence of different types of radio pulsar spectra owes to the environmental conditions around the neutron stars. Even assuming that the appearance of various non-standard spectra shapes in the general population of pulsars can be caused by peculiar environmental conditions, it is impossible to isolate different kinds of elements that influence pulsar radio emission and analysed them separately. However, we can notice a significant difference between typical broken and turnover spectrum.

4 SUMMARY

We can finalize that detailed analysis of the PSR B1259−63 spectra for observing days reveals that the shape of the spectrum depends on the orbital phase. Moreover, the observed evolution is widely consistent with changes of spectra averaged over orbital phase intervals. Furthermore, the fits presented in Tab.1 show significant differences between spectra before and after eclipse. The results leads us to conclusion that the PSR B1259−63 spectrum varies with the changes of the pulsar environment what can be explained by combination of two effects: free–free absorption and cyclotron resonance (Kijak et al. 2011a). At the same time the detailed analysis of the PSR B1259−63 spectra for observing days revealed appearance of all types of spectra.

Clearly, in the case of this pulsar, various spectra shapes are caused by the same physical phenomena, the differences in the shapes appear to be of an quantitative origin, rather than qualitative. We see the both the GPS type spectrum, and the broken type spectra (of various break “strengths”, ranging from the flat-normal spectra to almost regular power-law ones), apparently appearing due to the same effects, only giving a varying amount of absorption. Extending that to the study of the spectra of isolated pulsars, we believe that the differences between the various pulsar spectra shapes may arise from the amounts of certain effects present (like the absorption at low observing frequencies), rather than due to the existence of different physical phenomena. This would make the classification of the pulsar spectra in general purely morphological, a shape description only, and not a tool to discern between various phenomena causing the spectra to differ from the canonical power-law type. Combining this with the study of the GPS pulsars we can conclude, that it is most likely the pulsar environment that causes the spectra to differ from the standard power-law. In the extreme cases it is causing the appearance of a turnover in the spectrum, but in otherwise it is changing the spectra only slightly at lower frequencies, which results in the detection of a broken-type spectra.

Therefore we believe, that the case of B1259−63 can be treated as a key factor to our understanding of not only the GPS phenomenon, but also other types of untypical spectra as well.

ACKNOWLEDGMENTS

MD is a scholar within Sub-measure 8.2.2 Regional Innovation Strategies, Measure 8.2 Transfer of knowledge, Priority VIII Regional human resources for the economy Human Capital Operational Programme co-financed by European Social Fund and state budget.

REFERENCES

Connors T. W., Johnston S., Manchester R. N., McConnell D., 2002, MNRAS, 336, 1201
Johnston, S., Manchester, R.N., Lyne, A.G., Nicastro, L., Spyromilio, J., 1994, MNRAS, 268, 430
Johnston S., Manchester R. N., McConnell D., Campbell-Wilson D., 1999, MNRAS, 302, 277
Johnston S., Ball L., Wang N., Manchester R. N., 2005, MNRAS, 358, 1069
Kijak J.,Maron, O., 2004, in Camilo F., Gaensler B.M., eds, Proc. IAU Symp. 218, Young Neutron Stars and Their Environments. Astron. Soc. Pac., San Francisco, p. 339
Kijak J., Gupta Y., Krzeszowski K., 2007, A&A, 462, 699
Kijak, J., Dembska, M., Lewandowski, W., Melikidze, G., Sendyk, M., 2011a, MNRAS, 418, L114
Kijak, J., Lewandowski, W., Maron, O., Gupta, Y., Jessner, A., 2011b, A&A, 531, A16
Kong, S.W., Yu, Y.W., Huang, Y.F., Cheng, K.S, 2011, MNRAS, 416, 1067
Kuzmin A. D., Losovsky B. Ya., 2001, A&A, 368, 230
Lorimer D. R., Kramer M., 2005, Handbook of Pulsar Astronomy (Cambridge University Press)
Lorimer D. R., Yates J. A., Lyne A. G., Gould D. M., 1995, MNRAS, 273, 411
Malofeev V. M., Gil J., Jessner A., Malov I. F., Seiradakis J. H., Sieber W., Wielebinski R., 1994, A&A, 285, 201
Maron O., Kijak J., Kramer M., Wielebinski R., 2000, A&A S,147, 195
McClure-Griffiths, N. M., Johnston, S., Stinebring, D. R., Nicastro, L., 1998, ApJ, 492, L49
Negueruela, I., Rib, M., Herrero, A., Lorenzo, J., Khangulyan, D., Aharonian, F.A., 2011, ApJL, 732, L11
Sieber W., 1973, A&A S, 28, 237